

**VECTOR BUNDLES AND $SO(3)$ -INVARIANTS
FOR ELLIPTIC SURFACES II:
THE CASE OF EVEN FIBER DEGREE**

ROBERT FRIEDMAN

Introduction.

Let S be a simply connected elliptic surface with at most two multiple fibers. In this paper, the second in a series of three, we are concerned with describing moduli spaces of stable vector bundles V over S such that the restriction of $c_1(V)$ to a general fiber has the smallest possible nonzero degree, namely the product of the multiplicities, in the case where this product is even. We then apply this study toward a partial calculation of the corresponding Donaldson polynomial invariants of S . Our goal is the completion of the C^∞ classification of such surfaces, and the general outline of this classification has been described in the introduction to Part I. Aside from quoting a few results from Part I, this paper can however be read independently. On the other hand, the methods of this paper draw heavily on the book [4], and many arguments which are very similar to arguments in [4] are sketched or simply omitted. Roughly speaking, the new ingredients in the proof consist of the algebraic geometry of certain elliptic surfaces associated to S , which have a single multiple fiber of multiplicity two and are birational to double covers of rational ruled surfaces. The vector bundle parts of the argument run more or less parallel to the arguments in [4], with a few new cases to analyze.

The outline of this paper is as follows. In this paper, we shall only be concerned with elliptic surfaces S over $\mathbb{P}^1$ with multiple fibers of multiplicities $2m_1$ and m_2 , where m_2 is odd, and such that there exists a divisor Δ on S with $\Delta \cdot f = 2m_1m_2$, the minimum possible value, for a smooth fiber f . In this case, there is an associated surface $J^{m_1m_2}(S)$ defined in [3]. The surface $J^{m_1m_2}(S)$ fibers over $\mathbb{P}^1$ and the fiber over a point t lying under a smooth fiber f of S is $J^{m_1m_2}(f)$, the set of line bundles of degree m_1m_2 on the fiber f of S . The surface $J^{m_1m_2}(S)$ has an involution defined by $\lambda \in J^{m_1m_2}(f) \mapsto \mathcal{O}_f(\Delta|f) \otimes \lambda^{-1}$. The quotient of $J^{m_1m_2}(S)$ by this involution is birational to a rational ruled surface $\mathbb{F}_N$, and we describe the geometry of the double cover in detail. In Section 2, we describe the rough classification of stable bundles V on S with $c_1(V) = \Delta$. To each such bundle there is an associated bisection C of $J^{m_1m_2}(S)$ which is invariant under the involution, and so defines a section of the quotient ruled surface. In Section 3, we show that for general bundles V , V is determined up to finite ambiguity by the section of the ruled surface and the choice of a certain line bundle on the associated bisection C of $J^{m_1m_2}(S)$.

1991 *Mathematics Subject Classification*. Primary 14J60, 57R55; Secondary 14D20, 14F05, 14J27.

Research partially supported by NSF grant DMS-9203940

Thus, a Zariski open subset of the moduli space fibers over an open subset of the linear system of all sections on a certain rational ruled surface, and the fibers are a number of copies of the Jacobian $J(C)$ of the bisection C of $J^{m_1 m_2}(S)$. It is here that the asymmetry between $2m_1$ and m_2 becomes apparent: the number of connected components of the fiber is just m_2 . The reasons for this are explained following Lemma 2.4. Finally, in Section 4 we calculate the leading coefficient of a Donaldson polynomial invariant and show that it contains an “extra” factor of m_2 .

1. Geometry of elliptic surfaces.

We fix notation for this paper. Let $\pi: S \rightarrow \mathbb{P}^1$ be an elliptic surface with two multiple fibers F_{2m_1} and F_{m_2} of multiplicities $2m_1$ and m_2 , where $\gcd(2m_1, m_2) = 1$. We shall further assume that the reductions of the multiple fibers are smooth and that all other singular fibers are reduced and irreducible with just one ordinary double point. In other words, S is nodal in the terminology of [4]. Let $\kappa_S = \kappa \in H^2(S; \mathbb{Z})$ be the unique class such that $2m_1 m_2 \kappa = [f]$, where f is a general fiber of π . Finally we shall assume that there is a $2m_1 m_2$ -section Δ , i.e. a divisor Δ , not necessarily effective, with $\Delta \cdot f = 2m_1 m_2$, or equivalently $\Delta \cdot \kappa = 1$. By [4], there always exist such nodal elliptic surfaces. In particular S is algebraic.

We shall be concerned with the associated elliptic surface $J^{m_1 m_2}(S)$ defined in Section 2 of [3]. By the discussion in Section 2 of [3], $J^{m_1 m_2}(S)$ has exactly one multiple fiber of multiplicity two, above the point of $\mathbb{P}^1$ corresponding to F_{2m_1} . We shall denote this fiber by F_2 . Moreover, given the $2m_1 m_2$ -section Δ , there is an involution of $J^{m_1 m_2}(S)$ defined on the generic fiber S_η by $\lambda \mapsto \mathcal{O}_{S_\eta}(\Delta) \otimes \lambda^{-1}$. Since $J^{m_1 m_2}(S)$ is relatively minimal, this involution on the generic fiber extends to an involution on $J^{m_1 m_2}(S)$, which we shall denote by ι . The data of $J^{m_1 m_2}(S)$ and the involution ι do not depend on Δ , but only on the restriction of Δ to the generic fiber. A line bundle which has the same restriction as Δ to the generic fiber differs from Δ by a line bundle which is trivial on the generic fiber and thus is a multiple of κ . Up to twisting by a line bundle which is divisible by 2, the only possibilities are then Δ and $\Delta - \kappa$. Replacing Δ by $\Delta - \kappa$ replaces Δ^2 by $\Delta^2 - 2$ and thus changes $\Delta^2 \pmod{4}$. Of course $\Delta^2 \equiv 1 \pmod{2}$ since

$$\Delta \cdot K_S = 2m_1 m_2 (p_g + 1) - 2m_1 - m_2 \equiv 1 \pmod{2}.$$

Let us determine the fixed point set of ι .

Lemma 1.1. *The fixed point set of ι consists of a smooth 4-section together with two isolated fixed points on F_2 .*

Proof. The fixed point set of an involution on the smooth surface $J^{m_1 m_2}(S)$ must consist of a smooth curve together with some isolated fixed points. For a smooth nonmultiple fiber f , there are four divisors λ such that $2\lambda = f$. Thus there is a component of the fixed point set which is a 4-section of $J^{m_1 m_2}(S)$. Every other curve component of the fixed point set has trivial restriction to the generic fiber and (since the curve component is smooth) cannot meet the 4-section. Since all fibers are irreducible, there can be no other component, and the remaining fixed points are isolated.

Let us consider the possibilities for isolated fixed points away from F_2 . In an analytic neighborhood of a nonmultiple fiber, there is a section of the elliptic fibration. Using this section to make the local identification of $J^{m_1 m_2}(S)$ with $J^0(S)$, and

since the group of local sections is divisible, it is easy to see that, after a translation, we can assume that ι corresponds to the involution $x \mapsto -x$. Direct inspection shows that this involution has no isolated fixed points, even at the nodal fibers. To handle the multiple fiber, the explicit description of a neighborhood X of a multiple fiber shows that after a base change of order two, say $\tilde{X} \rightarrow X$, we can assume that there is a local section of $\tilde{X}$. The induced map from the central fiber of $\tilde{X}$ to the central fiber of X corresponds to taking the quotient by a subgroup of order two. After a translation we can further assume that the pulled back involution on $\tilde{X}$ is given by $x \mapsto -x$. Since inverses commute with translations by a point of order two, the restriction of ι to F_2 again has four fixed points. Two of these lie on the 4-section (recall that a 4-section can meet F_2 in at most two distinct points) and the remaining two are isolated. $\square$

Thus there are two isolated fixed points of ι on F_2 . If we blow these up and then take the quotient, the result will be a smooth surface $\mathbb{F}$ mapping to $\mathbb{P}^1$ whose fibers are smooth rational curves except over the point corresponding to F_2 (FIG. 1). Over this point, the fiber is a curve $\mathfrak{d}_1 + 2\mathfrak{e} + \mathfrak{d}_2$, where $\mathfrak{d}_1$ and $\mathfrak{d}_2$ are the images of the exceptional curves, $\mathfrak{e}$ is the image of F_2 , and we have $(\mathfrak{d}_i)^2 = -2$ and $\mathfrak{e}^2 = -1$, $\mathfrak{d}_1 \cdot \mathfrak{e} = \mathfrak{d}_2 \cdot \mathfrak{e} = 1$ and $\mathfrak{d}_1 \cdot \mathfrak{d}_2 = 0$. In particular we may contract $\mathfrak{e}$ and then either $\mathfrak{d}_1$ or $\mathfrak{d}_2$ to obtain a rational ruled surface $\mathbb{F}_N$. We shall fix notation so that $\mathfrak{d}_2$ is the curve we contract and the resulting surface is $\mathbb{F}_N$. However, as we shall see, it is important to keep in mind the symmetry between $\mathfrak{d}_1$ and $\mathfrak{d}_2$. Contracting $\mathfrak{d}_1$ instead corresponds to making an elementary modification of $\mathbb{F}_N$ and thus replacing it by $\mathbb{F}_{N \pm 1}$. As we shall see, the symmetry between $\mathfrak{d}_1$ and $\mathfrak{d}_2$ corresponds to the choice of either Δ or $\Delta - \kappa$.

The branch divisor B' on the blowup $\mathbb{F}$ of $\mathbb{F}_N$ is of the form $B + \mathfrak{d}_1 + \mathfrak{d}_2$, where B is a smooth 4-section which does not meet $\mathfrak{d}_1$ or $\mathfrak{d}_2$ and hence $B \cdot \mathfrak{e} = 2$. Thus if we use the basis $\{\sigma, f, \mathfrak{d}_2, \mathfrak{e}\}$ for $\text{Pic}(\mathbb{F})$, where σ is the negative section of $\mathbb{F}_N$ and f is the class of a fiber, viewed as curves on $\mathbb{F}$, it is easy to see that

$$B = 4\sigma + (2k + 1)f - 4\mathfrak{e} - 2\mathfrak{d}_2$$

for some odd integer k and that

$$B + \mathfrak{d}_1 + \mathfrak{d}_2 = 4\sigma + (2k + 2)f - 6\mathfrak{e} - 2\mathfrak{d}_2,$$

which is indeed divisible by 2. Note that we cannot say *a priori* that B is irreducible. However it cannot be a union of a 3-section and a section, since there are no sections of $J^{m_1 m_2}(S)$. In particular $B \cdot \sigma \geq 0$. Thus if $\mathfrak{d}_1$ is the proper transform of the fiber on $\mathbb{F}_N$, and we assume that the negative section σ does not pass through the point of the original fiber that was blown up, so that $\sigma \cdot \mathfrak{d}_2 = 0$, then $2k + 1 \geq 4N$, or equivalently

$$k \geq 2N.$$

The same conclusion holds by a similar argument if σ does pass through the point that is blown up.

It is now easy to reverse this procedure. Begin with $\mathbb{F}_N$ and blow up a point in a fiber. Then blow up the point of intersection of the exceptional curve with the proper transform of the fiber. The result is a nonminimal ruled surface $\mathbb{F}$ with a reducible fiber of the ruling of the form $\mathfrak{d}_1 + 2\mathfrak{e} + \mathfrak{d}_2$, where $\mathfrak{d}_1$ is the proper

transform of the fiber, $\mathfrak{d}_2$ is the proper transform of the first exceptional curve, and $\mathfrak{e}$ is the second exceptional curve. Choose a smooth element B in the linear system $|4\sigma + (2k+1)f - 4\mathfrak{e} - 2\mathfrak{d}_2|$, if any exist, where σ is the negative section of $\mathbb{F}_N$ and f is a fiber. The double cover of $\mathbb{F}$ branched along $B + \mathfrak{d}_1 + \mathfrak{d}_2$ is then an elliptic surface with a multiple fiber of multiplicity 2, bisections corresponding to the pullbacks of sections of $\mathbb{F}_N$, and an involution ι .

Let us calculate $p_g(J^{m_1 m_2}(S)) = p_g(S)$ in terms of $\mathbb{F}$ and B . The canonical bundle of $\mathbb{F}_N$ is given by $K_{\mathbb{F}_N} = -2\sigma - (N+2)f$. Thus, recalling that $\mathfrak{d}_2$ is the proper transform of the first exceptional curve and that $\mathfrak{e}$ is the second, we have

$$K_{\mathbb{F}} = -2\sigma - (N+2)f + \mathfrak{d}_2 + 2\mathfrak{e}.$$

As for the branch locus $B + \mathfrak{d}_1 + \mathfrak{d}_2$, we have

$$B + \mathfrak{d}_1 + \mathfrak{d}_2 = 2(2\sigma + (k+1)f - 3\mathfrak{e} - \mathfrak{d}_2).$$

By standard formulas for double covers,

$$H^0(J^{m_1 m_2}(S); K_{J^{m_1 m_2}(S)}) = H^0(\mathbb{F}; \mathcal{O}_{\mathbb{F}}(K_{\mathbb{F}} + 2\sigma + (k+1)f - 3\mathfrak{e} - \mathfrak{d}_2)).$$

Now, using the calculations above, we have

$$K_{\mathbb{F}} + 2\sigma + (k+1)f - 3\mathfrak{e} - \mathfrak{d}_2 = (k - N - 1)f - \mathfrak{e}.$$

Recalling that f is linearly equivalent to $\mathfrak{d}_1 + 2\mathfrak{e} + \mathfrak{d}_2$, it is clear that

$$h^0((k - N - 1)f - \mathfrak{e}) = \begin{cases} k - N - 1, & \text{if } k - N \geq 2 \\ 0, & \text{otherwise.} \end{cases}$$

Now if $k - N \leq 1$, since $k \geq 2N$ we must have $N \leq 1$. If $N = 1$, then $k = 2$ and so $k - N - 1 = 0$ in this case as well. If $N = 0$, then $k = 0$. In this case $B = 4\sigma + f - 4\mathfrak{e} - 2\mathfrak{d}_2 = 4\sigma + \mathfrak{d}_1 - 2\mathfrak{e} - \mathfrak{d}_2$, and it is easy to see that the effective curve $\sigma - \mathfrak{d}_2 - \mathfrak{e}$, which is the proper transform of the unique element of $|\sigma|$ passing through the point of the fiber which is blown up, satisfies

$$(\sigma - \mathfrak{d}_2 - \mathfrak{e}) \cdot (4\sigma + f - 4\mathfrak{e} - 2\mathfrak{d}_2) = -1.$$

Thus $|B|$ has the fixed component $\sigma - \mathfrak{d}_2 - \mathfrak{e}$, which is a section, and this case does not arise. So in all cases we have $h^0((k - N - 1)f - \mathfrak{e}) = k - N - 1$.

Thus we may summarize this discussion as follows:

Lemma 1.2. *With notation and conventions as above,*

$$p_g(S) = p_g(J^{m_1 m_2}(S)) = k - N - 1. \quad \square$$

In particular, suppose that $p_g(S) = 0$. Since $k \geq 2N$, and the case $k = N = 0$ has been ruled out above, the only possibilities are $N = 0, k = 1$ or $N = 1, k = 2$. Thus $\mathbb{F}$ is the blowup of either $\mathbb{F}_0$ or $\mathbb{F}_1$, and of course the two cases are elementary transformations of each other. Thus we may assume that $k = 2$ and $N = 1$ in this case. Moreover the negative section of $\mathbb{F}_1$ does not pass through the exceptional point in the blowup.

2. Classification of stable bundles.

We let S be a nodal elliptic over $\mathbb{P}^1$ with exactly two multiple fibers of multiplicities $2m_1$ and m_2 and let Δ be a divisor on S with $\Delta \cdot f = 2m_1m_2$. Fix an integer c . In this section we shall study rank two vector bundles V with $c_1(V) = \Delta$ and $c_2(V) = c$. We shall also let $w = \Delta \bmod 2$ and $p = \Delta^2 - 4c$.

First recall the following standard definition from [3]:

Definition 2.1. An ample line bundle L on S is (Δ, c) -suitable or (w, p) -suitable if for all divisors D on S such that $-D^2 + D \cdot \Delta \leq c$, either $f \cdot (2D - \Delta) = 0$ or

$$\text{sign } f \cdot (2D - \Delta) = \text{sign } L \cdot (2D - \Delta).$$

The following is Lemma 3.3 of [3]:

Lemma 2.2. For all pairs (Δ, c) , (Δ, c) -suitable ample line bundles exist. $\square$

With this said, here is the rough classification of rank two vector bundles V with $c_1(V) = \Delta$ and $c_2(V) = c$ which are stable with respect to a c -suitable line bundle L .

Theorem 2.3. Let L be (Δ, c) -suitable, and let V be an L -stable rank two vector bundle with $c_1(V) = \Delta$ and $c_2(V) = c$. Then there exist:

- (i) A smooth irreducible curve C and a birational map $C \rightarrow \overline{C} \subseteq J^{m_1m_2}(S)$, where $\overline{C}$ is a bisection of $J^{m_1m_2}(S)$ invariant under the involution ι ;
- (ii) A divisor D on T , the minimal desingularization of the normalization of $C \times_{\mathbb{P}^1} S$, such that $D \cdot f = m_1m_2$, where f is a general fiber of $T \rightarrow C$, and moreover D has the same restriction to the generic fiber of T as the divisor induced by the section of $J^{m_1m_2}(T)$ corresponding to the map $C \rightarrow J^{m_1m_2}(S)$;
- (iii) A codimension two local complete intersection Z and an exact sequence

$$0 \rightarrow \mathcal{O}_T(D) \rightarrow \nu^*V \rightarrow \mathcal{O}_T(\nu^*\Delta - D) \otimes I_Z \rightarrow 0,$$

where $\nu: T \rightarrow S$ is the natural degree two map.

Moreover the bisection $\overline{C}$ and the double cover T are uniquely determined by the bundle V , and D is determined by the bundle V and the choice of a map $C \rightarrow J^{m_1m_2}(S)$. Finally, every rank two vector bundle V with $c_1(V) = \Delta$ and $c_2(V) = c$ satisfying (i)–(iii) above is stable with respect to every (Δ, c) -suitable ample line bundle L .

Proof. First suppose that V is L -stable. It follows from Theorem 4.3 of [3] that the restriction of V to the geometric fiber of π is semistable. More precisely, let $\eta = \text{Spec } k(\mathbb{P}^1)$ and let $\bar{\eta} = \text{Spec } \overline{k(\mathbb{P}^1)}$, where $\overline{k(\mathbb{P}^1)}$ denotes the algebraic closure of $k(\mathbb{P}^1)$. Let $V_{\bar{\eta}}$ denote the pullback of V to the curve $S_{\bar{\eta}}$ which is the geometric fiber of π . Then $V_{\bar{\eta}}$ is semistable. By the classification of rank two bundles on an elliptic curve, $V_{\bar{\eta}} = L_1 \oplus L_2$, where each L_i is a line bundle over $S_{\bar{\eta}}$ of degree m_1m_2 and $L_1 \otimes L_2$ corresponds to the restriction of Δ to $S_{\bar{\eta}}$. The Galois group $\text{Gal}(\overline{k(\mathbb{P}^1)}/k(\mathbb{P}^1))$ permutes the set $\{L_1, L_2\}$. This action cannot be trivial, since otherwise L_i would be rational over $k(\mathbb{P}^1)$ and then S would have an m_1m_2 -section. Thus the fixed field of the subgroup of $\text{Gal}(\overline{k(\mathbb{P}^1)}/k(\mathbb{P}^1))$ which operates trivially on $\{L_1, L_2\}$ defines a

degree two extension of $k(\mathbb{P}^1)$, corresponding to a morphism $C \rightarrow \mathbb{P}^1$. Setting T to be the minimal resolution of the normalization of $C \times_{\mathbb{P}^1} S$, there is a section of $J^{m_1 m_2}(T)$ defined by L_1 , say. The image of this section in $J^{m_1 m_2}(S)$ is then the bisection $\overline{C}$. By construction $\overline{C}$ is invariant under the involution ι . Let $\nu: T \rightarrow S$ be the natural degree two map.

The inclusion $L_1 \rightarrow V_{\overline{\eta}}$ induces a sub-line bundle $\mathcal{O}_T(D) \rightarrow \nu^*V$, which we may assume to have torsion free cokernel. Since $L_2 \neq L_1$, it is clear that this sub-line bundle is unique. The quotient is then necessarily of the form $\mathcal{O}_T(\nu^*\Delta - D) \otimes I_Z$.

It remains to prove that every V satisfying the above description is indeed L -stable. It follows from (iii) that $V_{\overline{\eta}}$ is an extension of two line bundles of degree $m_1 m_2$ and is therefore semistable. Again using Theorem 4.3 of [3], V is L -stable. $\square$

Next we discuss the meaning of the scheme Z and the bisection $\overline{C}$. The following is the analogue of Lemma 1.11 in [4] and is proved in exactly the same way:

Lemma 2.4. *Let f be a smooth fiber of π and let g be a component of $\nu^{-1}(f)$. Then $\text{Supp } Z \cap g \neq \emptyset$ if and only if $V|_f$ is unstable. In particular, if ν is not branched over f , so that $\nu^{-1}(f) = g \cup g'$, then $\text{Supp } Z \cap g \neq \emptyset$ if and only if $\text{Supp } Z \cap g' \neq \emptyset$. $\square$*

Next we turn to the section $\overline{C}$. Since $\overline{C}$ is invariant under ι , its proper transform on the blowup of $J^{m_1 m_2}(S)$ at the two isolated fixed points of ι is the pullback of a section A' of $\mathbb{F}$, which in turn induces a section A of $\mathbb{F}_N$. We shall use throughout the notation and conventions of the previous section. Notice that the section A' meets the reducible fiber either along $\mathfrak{d}_1$ or $\mathfrak{d}_2$, the two components of multiplicity one. Here $A' \cdot \mathfrak{d}_1 = 1$ and $A' \cdot \mathfrak{d}_2 = 0$ if A does not pass through the point of the corresponding fiber of $\mathbb{F}_N$ which was blown up, and $A' \cdot \mathfrak{d}_2 = 1$ and $A' \cdot \mathfrak{d}_1 = 0$ in the remaining case. Since the branch locus of the map $J^{m_1 m_2}(S) \rightarrow \mathbb{F}$ consists of $B + \mathfrak{d}_1 + \mathfrak{d}_2$, we see that A' always passes through the branch locus over the point corresponding to F_2 . Of course, this is also clear from the picture on $J^{m_1 m_2}(S)$: since $\overline{C}$ is a bisection, $\overline{C} \cdot f = 2$ and therefore $\overline{C} \cdot F_2 = 1$. It follows that $\overline{C}$ is smooth at the point of intersection with F_2 , that the intersection is transverse, and that the natural map $\overline{C} \rightarrow \mathbb{P}^1$ is always branched at the point corresponding to F_2 . A similar statement will hold for the map $C \rightarrow \mathbb{P}^1$. This fact is the fundamental difference between the case studied in this paper and the case of trivial determinant studied in [2] and [4].

Since A' always meets $\mathfrak{d}_1$ or $\mathfrak{d}_2$, it follows that the inverse image of A' in the blowup of $J^{m_1 m_2}(S)$ always meets exactly one of the two exceptional curves, and in fact meets it transversally at one point. Thus the inverse image of A' is the proper transform of $\overline{C}$, and therefore

$$(\overline{C})^2 = 2(A')^2 + 1.$$

Let us consider the section A of $\mathbb{F}_N$ in more detail. Either $A = \sigma$ or $A \in |\sigma + (N + s)f|$ for a uniquely specified nonnegative integer f . Moreover either A does not pass through the point on $\mathbb{F}_N$ which is the image of the exceptional divisor, in which case $A' \cdot \mathfrak{d}_2 = 0$, or it does, in which case $A' \cdot \mathfrak{d}_2 = 1$. The following lemma relates the odd integer $\Delta^2 - 4c = p$ to the invariants of V :

Lemma 2.5. *With notation as above, denote by the exceptional point the point of $\mathbb{F}_N$ which is blown up under the morphism $\mathbb{F} \rightarrow \mathbb{F}_N$. Then, if we set $p = p_1(\text{ad } V) =$*

$$\Delta^2 - 4c,$$

$$-p = \begin{cases} 4k - 6N + 1 + 2\ell(Z) + \delta, & \text{if } A = \sigma \text{ does not pass through} \\ & \text{the exceptional point;} \\ 4s + 4k - 2N + 1 + 2\ell(Z) + \delta, & \text{if } A \in |\sigma + (N + s)f| \text{ does not pass} \\ & \text{through the exceptional point;} \\ 4k - 6N - 1 + 2\ell(Z) + \delta, & \text{if } A = \sigma \text{ passes through} \\ & \text{the exceptional point;} \\ 4s + 4k - 2N - 1 + 2\ell(Z) + \delta, & \text{if } A \in |\sigma + (N + s)f| \text{ passes through} \\ & \text{the exceptional point.} \end{cases}$$

Here δ is a nonnegative integer which is zero if the map $C \rightarrow \mathbb{P}^1$ is not branched over any point corresponding to a singular nonmultiple fiber of $\pi: S \rightarrow \mathbb{P}^1$.

Proof. Since $c_1^2(\nu^*V) = 2c_1^2(V) = 2\Delta^2$ and $c_2(\nu^*V) = 2c_2(V) = 2c$, it will suffice to work with ν^*V . Clearly

$$c_2(\nu^*V) = -D^2 + D \cdot \nu^*\Delta + \ell(Z).$$

Thus

$$2(4c - \Delta^2) = -(2D - \nu^*\Delta)^2 + 4\ell(Z).$$

Now we can write

$$2D - \nu^*\Delta = D - (\nu^*\Delta - D),$$

where both D and $\nu^*\Delta - D$ naturally correspond to sections of $J^{m_1 m_2}(T)$. In fact, if $\varphi: J^{m_1 m_2}(T) \rightarrow J^{m_1 m_2}(S)$ is the obvious map, then the bisection $\overline{C}$ satisfies $\varphi^*\overline{C} = C_1 + C_2$, where C_1 and C_2 are sections of $J^{m_1 m_2}(T)$ corresponding to the divisors D and $\nu^*\Delta - D$ on T . An argument essentially identical to the proofs of Claim 1.17 and 1.18 in Chapter 7 of [4] shows that there is a nonnegative integer δ such that

$$-(2D - \nu^*\Delta)^2 = -(C_1 - C_2)^2 + 2\delta.$$

Moreover $\delta = 0$ if the map $C \rightarrow \mathbb{P}^1$ is not branched over any point corresponding to a singular nonmultiple fiber of $\pi: S \rightarrow \mathbb{P}^1$. Thus we must calculate $(C_1 - C_2)^2$. But, using the fact that C_i is a section of $J^{m_1 m_2}(T)$, we have

$$(C_i)^2 = -2(1 + p_g(S)) = -2(k - N).$$

Moreover $C_1 + C_2 = \varphi^*\overline{C}$. Thus

$$\begin{aligned} (C_1 - C_2)^2 &= 2(C_1)^2 + 2(C_2)^2 - (C_1 + C_2)^2 \\ &= -8(k - N) - 2(\overline{C})^2 \\ &= -8(k - N) - 4(A')^2 - 2. \end{aligned}$$

Clearly we have

$$(A')^2 = \begin{cases} -N, & \text{if } A = \sigma \text{ does not pass through the exceptional point;} \\ N + 2s, & \text{if } A \in |\sigma + (N + s)f| \text{ does not pass through} \\ & \text{the exceptional point;} \\ -N - 1, & \text{if } A = \sigma \text{ passes through the exceptional point;} \\ N + 2s - 1, & \text{if } A \in |\sigma + (N + s)f| \text{ passes through} \\ & \text{the exceptional point.} \end{cases}$$

Putting these formulas together gives the statement of the lemma. $\square$

Using Lemma 2.5, the inequality $k \geq 2N$ and the fact that if $N = 0$ then $k \geq 1$, whereas if $N = 1$ and $k = 2$ then the section σ does not pass through the exceptional point, we can easily deduce the following slight strengthening of Bogomolov's inequality in our case:

Corollary 2.6. *We have the following inequality for $-p$:*

$$-p \geq \begin{cases} 4p_g(S) - 2N + 5 & \text{if } A = \sigma \text{ does not pass through the exceptional point;} \\ 4p_g(S) + 2N + 5 & \text{if } A \in |\sigma + (N + s)f| \text{ does not pass through} \\ & \text{the exceptional point;} \\ 4p_g(S) - 2N + 3 & \text{if } A = \sigma \text{ passes through the exceptional point;} \\ 4p_g(S) + 2N + 3 & \text{if } A \in |\sigma + (N + s)f| \text{ passes through} \\ & \text{the exceptional point.} \end{cases}$$

In all cases $-p \geq 2p_g(S) + 1$, and if $p_g(S) = 0$, then $-p \geq 3$. $\square$

3. A Zariski open subset of the moduli space.

Our goal in this section is to prove the following theorem:

Theorem 3.1. *Let V be an L -stable rank two bundle on S . Suppose that*

- (i) *The associated bisection $\overline{C}$ of $J^{m_1 m_2}(S)$ is smooth, or equivalently that $\overline{C} = C$, and the image of $\overline{C}$ in $\mathbb{F}$ is not the proper transform of σ ;*
- (ii) *The map $C \rightarrow \mathbb{P}^1$ is not branched at any point corresponding to a singular fiber of π or at the multiple fiber of odd multiplicity m_2 ;*
- (iii) *The scheme Z on the associated double cover T is empty, and thus there is an exact sequence*

$$0 \rightarrow \mathcal{O}_T(D) \rightarrow \nu^*V \rightarrow \mathcal{O}_T(\nu^*\Delta - D) \rightarrow 0.$$

Then $V = \nu_\mathcal{O}_T(D + F) = \nu_*\mathcal{O}_T(\nu^*\Delta - D)$. In particular V is uniquely determined by the associated section A of $\mathbb{F}_N$ and the divisor D on T . Finally V is a smooth point of its moduli space, which is of dimension $-p - 3\chi(\mathcal{O}_S)$ at V .*

It is clear that the conditions above are equivalent to assuming that A' meets the branch locus B transversally, and that no point of intersection lies over a point of $\mathbb{P}^1$ corresponding to a singular nonmultiple fiber or to the multiple fiber of multiplicity m_2 , and that $Z = \emptyset$.

The proof of (3.1) will proceed along lines very similar to the proof of Theorem 1.12 in Chapter 7 of [4], and we shall simply sketch some of the details.

Let A be the section of $\mathbb{F}_N$ corresponding to A' . By assumption $A \neq \sigma$. Let r be the nonnegative integer such that $A \in |\sigma + (N + r)f|$. If the section A does not pass through the exceptional point of the blowup, then

$$(A') \cdot (B + \mathfrak{d}_1 + \mathfrak{d}_2) = (\sigma + (N + r)f) \cdot (4\sigma + (2k + 2)f - 6\mathfrak{e} - 2\mathfrak{d}_2) = 4r + 2k + 2.$$

Of these points, one corresponds to the intersection $A' \cdot \mathfrak{d}_1$, and so the branch divisor of the map $T \rightarrow S$ is $(4r + 2k + 1)f$, where f is a general fiber of π . This divisor is

even since f is divisible by 2, and we set $G = (4r + 2k + 1)f/2$. Likewise, if A does pass through the exceptional point of the blowup, then

$$(A') \cdot (B + \mathfrak{d}_1 + \mathfrak{d}_2) = (\sigma + (N + r)f - \mathfrak{d}_2 - \mathfrak{e}) \cdot (4\sigma + (2k + 2)f - 6\mathfrak{e} - 2\mathfrak{d}_2) = 4r + 2k.$$

In this case we set $G = (4r + 2k - 1)f/2$. Let F be the branch divisor in T , so that $\nu^*G \equiv F$. Thus $F = (4r + 2k + 1)f$ or $(4r + 2k - 1)f$. For future reference, let us also record the genus of C :

Lemma 3.2. *Let C satisfy (i) and (ii) of (3.1). Then*

$$g(C) = \begin{cases} 2r + k, & \text{if } A \text{ does not pass through the exceptional point;} \\ 2r + k - 1, & \text{if } A \text{ passes through the exceptional point.} \end{cases}$$

Proof. The map $C \rightarrow \mathbb{P}^1$ is branched at $A' \cdot (B + \mathfrak{d}_1 + \mathfrak{d}_2) = 4r + 2k + 2$ points if A does not pass through the exceptional point, and $4r + 2k$ points otherwise. The lemma now follows from the Riemann-Hurwitz formula. $\square$

Now $\det \nu_* \mathcal{O}_T(D) = \nu_* D - G$. Clearly $\nu_* D$ and Δ have the same restriction to the generic fiber. Arguing as in Chapter 7, (1.20) of [4], there is an injective map

$$\nu_* \mathcal{O}_T(D + F) \rightarrow V.$$

Set $W = \nu_* \mathcal{O}_T(D + F)$. Our goal will be to show that $W = V$. Preliminary to this goal we shall analyze W and the map $W \rightarrow V$. As a divisor class $\det W = \nu_* D - G + 2G = \nu_* D + G$. In addition there is an effective divisor E such that $(\det W)^{-1} \otimes \det V = \mathcal{O}_S(E)$. Thus $\det W = \Delta - E$ and E has trivial restriction to the generic fiber, so that E is a union of fibers (possibly including the reductions of the multiple fibers). Moreover

$$\nu_* D = \Delta - G - E.$$

Set $E' = \nu^* E$. We have

$$D + \iota^* D = \nu^* \nu_* D = \nu^* \Delta - F - E',$$

and therefore

$$\iota^* D = \nu^* \Delta - D - F - E'.$$

We can thus write

$$W = \nu_* \mathcal{O}_T(D + F) = \nu_* \mathcal{O}_T(\nu^* \Delta - D - E').$$

Using the fact that there is a surjection from $\nu^* W$ to $\mathcal{O}_T(\nu^* \Delta - D - E')$, we conclude that there is an exact sequence

$$0 \rightarrow \mathcal{O}_T(D) \rightarrow \nu^* W \rightarrow \mathcal{O}_T(\nu^* \Delta - D - E') \rightarrow 0.$$

Comparing this sequence to the defining exact sequence for $\nu^* V$ and arguing as in (1.24) of Chapter 7 of [4], we may conclude:

Lemma 3.3. *Let $Q = V/W$. Then $\nu^*Q \cong [\mathcal{O}_T/\mathcal{O}_T(-E')] \otimes \mathcal{O}_T(\nu^*\Delta - D)$. $\square$*

Our goal now is to prove the following:

Lemma 3.4. *In the above notation, $E' = 0$. Thus $Q = 0$ and $V = W = \nu_*\mathcal{O}_T(D + F)$ where $\nu_*D = \Delta - G$.*

We begin with the following construction. Let e be a component of the support of E' , and write $E' = ae + E''$, where E'' is effective and disjoint from e and $a > 0$. If e is not the multiple fiber of multiplicity m_1 on T , then either ν is unbranched over e or e is a smooth fiber. In either of these cases ν induces an isomorphism from e to $\nu(e) = f$, and we shall identify $\nu(e)$ with e . In the remaining case $e = F_{m_1}$ is the multiple fiber of multiplicity m_1 and ν is an étale double cover. There is then the following analogue of (1.25) of Chapter 7 of [4]:

Lemma 3.5. *There is a subsheaf Q_0 of ν_*Q which is isomorphic to*

- (i) $\mathcal{O}_e(-(a-1)e + \nu^*\Delta - D)$, viewed as a sheaf on $\nu(e) = f$, if $e \neq F_{m_1}$;
- (ii) A line bundle on F_{2m_1} such that

$$\nu^*Q_0 \cong \mathcal{O}_{F_{m_1}}(-(a-1)F_{m_1} + \nu^*\Delta - D),$$

in case $e = F_{m_1}$.

Proof. The argument in case $e \neq F_{m_1}$ runs as in (1.25) of Chapter 7 of [4]. If $e = F_{m_1}$, then ν^*Q contains the subsheaf

$$Q'_0 = [\mathcal{O}_T(-(a-1)e/\mathcal{O}_T(-ae))] \otimes \mathcal{O}_T(\nu^*\Delta - D),$$

which is a line bundle on F_{m_1} . The vector bundle $\nu_*Q'_0$ is a rank two vector bundle on F_{2m_1} with $\deg(\nu_*Q'_0) = m_1m_2$. Consider the rank two vector bundle $\nu^*\nu_*Q'_0$ on F_{m_1} . Its determinant is $Q'_0 \otimes \iota^*Q'_0 = (Q'_0)^{\otimes 2}$ (recall that $\nu^*\Delta - D$ is fixed under the involution) and there is a surjective map $\nu^*\nu_*Q'_0 \rightarrow Q'_0$. Thus there is an exact sequence

$$0 \rightarrow Q'_0 \rightarrow \nu^*\nu_*Q'_0 \rightarrow Q'_0 \rightarrow 0.$$

It follows that $\nu^*\nu_*Q'_0$ is semistable and is either $Q'_0 \oplus Q'_0$ or the nontrivial extension of Q'_0 by Q'_0 . On the other hand, $\nu_*Q'_0$ is either a direct sum of line bundles, say $Q_0 \oplus Q_1$ for two line bundles Q_i or a nontrivial extension of a line bundle Q_0 by Q_0 . In the first case we must have $\nu^*Q_i = Q'_0$ and in the second $\nu^*Q_0 = Q'_0$. In either case there is a subbundle Q_0 of ν_*Q as desired. $\square$

To prove Lemma 3.4, we shall assume that $E' \neq 0$ and derive a contradiction. Again, the argument will be very similar to the argument given in [4]. It will suffice to show that $\dim \text{Ext}^1(Q_0, W) \leq 1$. We also have that

$$\text{Ext}^1(Q_0, W) = H^0(W \otimes \mathcal{O}_S(e) \otimes Q_0^{-1}).$$

The case where e does not lie over the branch locus of $C \rightarrow \mathbb{P}^1$ follows exactly as in [4]. The case where e is a smooth fiber in the branch locus also follows by these methods provided we can show that $R^1\rho_*\mathcal{O}_T(2D - \nu^*\Delta)$ has length one at the point of C corresponding to e . This is a local calculation, which we shall leave to the

reader; it uses the fact that A' meets the branch locus transversally and can in fact be deduced from the global argument in [4], proof of Lemma 1.19 of Chapter 7.

There remains the new case, where $e = F_{m_1}$. In this case, the natural map

$$W = \nu_* \mathcal{O}_T(\nu^* \Delta - D - E') \rightarrow \nu_* \mathcal{O}_{F_{m_1}}(\nu^* \Delta - D - aF_{m_1})$$

is surjective, as one can see from applying the surjective map ν_* to the exact sequence

$$0 \rightarrow \mathcal{O}_T(\nu^* \Delta - D - E' - F_{m_1}) \rightarrow \mathcal{O}_T(\nu^* \Delta - D - E') \rightarrow \mathcal{O}_{F_{m_1}}(\nu^* \Delta - D - aF_{m_1}) \rightarrow 0.$$

It follows that $W|_{F_{2m_1}} = \nu_* \mathcal{O}_{F_{m_1}}(\nu^* \Delta - D - aF_{m_1})$. Thus

$$\begin{aligned} H^0(W \otimes \mathcal{O}_S(F_{2m_1}) \otimes Q_0^{-1}) &= H^0(\nu_* \mathcal{O}_{F_{m_1}}(\nu^* \Delta - D - aF_{m_1}) \otimes \mathcal{O}_S(F_{2m_1}) \otimes Q_0^{-1}) \\ &= H^0(\nu_* [\mathcal{O}_{F_{m_1}}(\nu^* \Delta - D - aF_{m_1}) \otimes \nu^* \mathcal{O}_S(F_{2m_1}) \otimes \nu^* Q_0^{-1}]) = H^0(\mathcal{O}_{F_{m_1}}), \end{aligned}$$

where we have used the fact that $\nu^* Q_0 = \mathcal{O}_{F_{m_1}}(\nu^* \Delta - D - (a-1)F_{m_1})$. Hence

$$\dim \text{Ext}^1(Q_0, W) = h^0(W \otimes \mathcal{O}_S(F_{2m_1}) \otimes Q_0^{-1}) = 1$$

as desired. $\square$

We see that we have proved all of Theorem 3.1 except the statement about the smoothness of the moduli space, which follows from:

Lemma 3.6. *Suppose that $V = \nu_* \mathcal{O}_T(D + F)$ as above. Then V is good, in the terminology of [2]. In other words, $H^2(S; \text{ad } V) = 0$.*

Proof. It suffices to show that $\dim \text{Hom}(V, V \otimes K_S) = h^0(K_S)$. Now

$$\begin{aligned} \text{Hom}(V, V \otimes K_S) &= \text{Hom}(V, \nu_* (\mathcal{O}_T(\nu^* \Delta - D) \otimes \nu^* K_S)) \\ &= \text{Hom}(\nu^* V, \mathcal{O}_T(\nu^* \Delta - D) \otimes \nu^* K_S). \end{aligned}$$

Using the defining exact sequence for ν^* , there is an exact sequence

$$0 \rightarrow H^0(\nu^* K_S) \rightarrow \text{Hom}(\nu^* V, \mathcal{O}_T(\nu^* \Delta - D) \otimes \nu^* K_S) \rightarrow H^0(\mathcal{O}_T(\nu^* \Delta - 2D) \otimes \nu^* K_S).$$

Since K_S is a rational multiple of the fiber and $\nu^* \Delta - 2D$ is nontrivial on the generic fiber, the term $H^0(\mathcal{O}_T(\nu^* \Delta - 2D) \otimes \nu^* K_S)$ is zero. Thus $\dim \text{Hom}(V, V \otimes K_S) = h^0(\nu^* K_S)$. Using the isomorphism $H^0(\nu^* K_S) \cong H^0(K_S) \oplus H^0(K_S(-G))$, it suffices to show that $H^0(K_S(-G)) = 0$. Now $K_S = \mathcal{O}_S((k - N - 2)f + (2m_1 - 1)F_{2m_1} + (m_2 - 1)F_{m_2})$. Also $G = (2r + k \pm 1/2)f$. Thus it suffices to observe that the linear system

$$|(-N - 2r - 2 \pm 1/2)f + (2m_1 - 1)F_{2m_1} + (m_2 - 1)F_{m_2}|$$

is empty. $\square$

Next let us describe the subset of the moduli space consisting of bundles V which satisfy the hypotheses of Theorem 3.1. We begin by reversing the procedure outlined above. Fix the section A' , which is generic in the sense of Theorem 3.1: it

meets B transversally and no point of intersection corresponds to a singular non-multiple fiber or to the multiple fiber of odd multiplicity. The section A' determines the bisection $C = \overline{C}$, and thus a double cover $\nu: T \rightarrow S$ together with an elliptic fibration $\rho: T \rightarrow C$ and a divisor D_0 , well-defined on the generic fiber. Moreover by construction $\nu_* D_0$ and Δ have the same restriction on the generic fiber, and thus differ by a multiple of κ . It is easy to see that changing D_0 by a sum of fiber components on T replaces $\nu_* D_0$ by an arbitrary even multiple of κ . Thus we may assume that we have $\nu_* D_0 = \Delta - G$ or $\nu_* D_0 = \Delta - \kappa - G$. It is an exercise in the formulas of the preceding section to see that we have

$$\Delta^2 \equiv 2(A')^2 + 1 \pmod{4}.$$

Additionally

$$2(A')^2 + 1 \equiv \begin{cases} 2N + 1 \pmod{4}, & \text{if } A \text{ does not pass through the exceptional point;} \\ 2N - 1 \pmod{4}, & \text{otherwise.} \end{cases}$$

Thus the symmetry between the possibility that A does or does not pass through the exceptional point, which is essentially the choice of blowing $\mathbb{F}$ down to $\mathbb{F}_N$ or $\mathbb{F}_{N \pm 1}$, reflects the choice of Δ or $\Delta - \kappa$, which in turn reflects $\Delta^2 \pmod{2}$, or equivalently the dimension of the moduli space $\pmod{2}$.

Having made one choice for a line bundle D_0 on the double cover T , where D_0 is specified on the generic fiber of T and satisfies $\nu_* D_0 = \Delta - G$, or $\nu_* D_0 = \Delta - \kappa - G$, the possibilities for D are given by the next lemma.

Lemma 3.7. *Given the double cover $T \rightarrow S$, the set of all D whose restriction to the generic fiber equals D_0 and which satisfy $\nu_* D = \Delta - G$, or $\nu_* D = \Delta - \kappa - G$ is a principal homogeneous space over $\text{Pic}^T T$, which in turn is an extension of the Jacobian $J(C)$ by a cyclic group of order m_2 . Moreover this principal homogeneous space is nonempty for exactly one of the two choices for $\nu_* D$ above.*

Proof. By the remarks preceding the lemma, there exists a D_0 with $\nu_* D_0 = \Delta - G$, or $\nu_* D_0 = \Delta - \kappa - G$, and only one of these possibilities can hold. If D has the same restriction to the generic fiber as D_0 and $\nu_* D = \nu_* D_0$, then $D - D_0$ has trivial restriction to the generic fiber and $\nu_*(D - D_0) = 0$. The first condition says that $D - D_0$ is of the form $\rho^* \lambda \otimes \mathcal{O}_T(aF_{m_1} + bF'_{m_2} + cF''_{m_2})$, where F_{m_1} is the multiple fiber of multiplicity m_1 lying above F_{2m_1} and F'_{m_2}, F''_{m_2} are the two multiple fibers of multiplicity m_2 lying over F_{m_2} . We may further assume that $0 \leq a < m_1$ and that $0 \leq b < m_2, 0 \leq c < m_2$. Here λ is a line bundle of degree d on C . Thus

$$\nu_*(D - D_0) = df + 2aF_{2m_1} + (b + c)F_{m_2}.$$

It is easy to see that this divisor is trivial if and only if $d = 0$, $a = 0$, and $b \equiv -c \pmod{m_2}$. Thus, there is a natural identification of the set of all D (given the fixed divisor D_0) with $J(C) \times \mathbb{Z}/m_2\mathbb{Z}$. $\square$

We now assume that A is not the negative section of $\mathbb{F}_N$ and write $A \in |\sigma + (N + r)f|$ where $r \geq 0$. The dimension of the linear system A' is then equal to n , where

$$n = \begin{cases} N + 2r + 1, & \text{if } A \text{ does not pass through the exceptional point;} \\ N + 2r, & \text{otherwise.} \end{cases}$$

We also let $g = g(C)$ be the genus of the bisection C , as given in Lemma 3.2. Note that

$$n + g = \begin{cases} 4r + N + k + 1 & \text{if } A \text{ does not pass through the exceptional point;} \\ 4r + N + k - 1 & \text{otherwise.} \end{cases}$$

In both cases, comparing this with the formula for $-p$ given in Lemma 2.5 and using the fact that $1 + p_g(S) = k - N$, we see that

$$-p - 3\chi(\mathcal{O}_S) = g + n.$$

Note finally that the moduli space will be nonempty provided that $-p \geq 4p_g(S) + 2N + 3$. Since $p_g(S) = k - N - 1 \geq N - 1$, the moduli space will be nonempty as long as

$$-p \geq 6p_g(S) + 5.$$

Arguing as in Theorem 1.14 of Chapter 7 of [4], we obtain the following:

Theorem 3.8. *Let p be an odd negative integer, and choose $w \in H^2(S; \mathbb{Z}/2\mathbb{Z})$ such that $w = \Delta \bmod 2$ or $w = \Delta - \kappa \bmod 2$, and that $w^2 \equiv p \bmod 4$. Let L be a (w, p) -suitable ample line bundle on S . Let $\mathfrak{M} = \mathfrak{M}(S, L; w, p)$ denote the moduli space of L -stable rank two bundles on S with $w_2(V) = w$ and $p_1(\text{ad } V) = p$. Then for all p such that $-p \geq 6p_g(S) + 5$, $\mathfrak{M}$ contains a nonempty Zariski open subset M corresponding to vector bundles V satisfying the hypotheses of Theorem 3.1. The set M is smooth of dimension $-p - 3\chi(\mathcal{O}_S)$. Moreover, there is a holomorphic map from M to a Zariski open subset $U \subseteq \mathbb{P}^n$ and the fibers are isomorphic to m_2 copies of a complex torus of dimension g . $\square$*

Let us finally consider the case where $p_g(S) = 0$ and $-p = 3$, the case of a moduli space of expected dimension zero. In this case we fix $N = 1$ and $k = 2$, and the negative section does not pass through the exceptional point. The Chern class calculations of Lemma 2.5 show that we must have $\ell(Z) = \delta = 0$ and A must be the negative section σ of $\mathbb{F}_1$. Note that $\sigma \cdot (B + \mathfrak{d}_1 + \mathfrak{d}_2) = 2$, and the intersection must be transverse since σ cannot split into a union of two sections in the double cover. Thus $C = \overline{C} \rightarrow \mathbb{P}^1$ is branched at two points, so that $C = \mathbb{P}^1$ again. Assuming as we may that the multiple fiber of odd multiplicity does not correspond to a branch point, we see that the arguments of Theorem 3.1 go through to show that there are exactly m_2 vector bundles V whose associated section is σ . (Here, in case the intersection point of σ with B corresponds to a singular nonmultiple fiber, we must use the more detailed analysis of [2] (5.12) and (5.13) to see that the section σ and the line bundle on T determine V .) Each of these is a smooth point of the moduli space, by a slight modification of the proof of Lemma 3.6 (in this case $G = f/2$ and $K_S = \mathcal{O}_S(-f + (2m_1 - 1)F_{2m_1} + (m_2 - 1)F_{m_2})$). Summarizing, then:

Theorem 3.9. *In case $p_g(S) = 0$, the moduli space corresponding to $-p = 3$ consists of m_2 reduced points. $\square$*

4. Calculation of the leading coefficient.

Fix w and p with $w^2 \equiv p \bmod 4$, and let $\mathfrak{M} = \mathfrak{M}(S, L; w, p)$ denote the moduli space of L -stable rank two bundles on S with $w_2(V) = w$ and $p_1(\text{ad } V) = p$. Let d be the (complex) dimension of $\mathfrak{M}$:

$$d = \begin{cases} 4r + N + k + 1 & \text{if } A \text{ does not pass through the exceptional point} \\ 4r + N + k - 1 & \text{otherwise,} \end{cases}$$

where the section $A \in |\sigma + (N+r)f|$. With this notation we see that $2n = d - p_g(S)$ and that $g = d - n$. Finally let us recall from (3.4) of [3] that in case $p_g(S) = 0$ there is a unique chamber of type (w, p) which contains κ in its closure, called the (w, p) -suitable chamber. We can now state the main result of this paper:

Theorem 4.1. *For $p_g(S) > 0$, let $\gamma_{w,p}(S, \beta)$ be the Donaldson polynomial corresponding to the $SO(3)$ -bundle P over S with $w_2(P) = \Delta \bmod 2$ and $p_1(P) = p$, and β is a choice of orientation agreeing with the usual complex orientation for $\mathfrak{M}$. If $p_g(S) = 0$, let $\gamma_{w,p}(S, \beta)$ be the corresponding Donaldson polynomial for metrics whose associated self-dual harmonic 2-form lies in the (w, p) -suitable chamber. Then, writing $\gamma_{w,p}(S, \beta)$ as a polynomial in κ_S and q_S , say $\gamma_{w,p}(S, \beta) = \sum_{i=0}^{[d/2]} a_i q_S^i \kappa_S^{d-2i}$, we have, for all p with $-p \geq 2(4p_g + 2)$, $a_i = 0$ for $i > n$ and*

$$a_n = \frac{d!}{2^n n!} (2m_1 m_2)^{p_g(S)} m_2.$$

We first remark that the assumption that $-p \geq 2(4p_g + 2)$ implies that $\mathfrak{M}$ is nonempty and contains a smooth Zariski open subset as described in Theorem 3.8. Indeed $-p$ is an odd integer greater than $8p_g + 4$, and so $-p \geq 8p_g + 5 \geq 6p_g + 5$. Thus we are in the range of Theorem 3.8.

Let $X = X_{w,p}$ denote the Uhlenbeck compactification associated to $\mathfrak{M}$ [1], [4]. The orientation of $\mathfrak{M}$ induces a fundamental class of X . There is a μ -map $H_2(S) \rightarrow H^2(\mathfrak{M})$, which roughly speaking is given by taking slant product with the class $-p_1(P)/4$, where P is the universal $SO(3)$ -bundle over $S \times \mathfrak{M}$. If P lifts to a holomorphic bundle $\mathcal{V}$ over $S \times \mathfrak{M}$, then $p_1(P) = c_1^2(\mathcal{V}) - 4c_2(\mathcal{V}) = p_1(\text{ad } \mathcal{V})$. The classes $\mu(\alpha) \in H^2(\mathfrak{M})$ extend uniquely to classes in $H^2(X)$, which we shall also denote by $\mu(\alpha)$. The Donaldson polynomial is then defined by taking cup products of the $\mu(\alpha)$ and evaluating on the fundamental cycle of X .

Arguing as in [4], it is a combinatorial exercise to deduce Theorem 4.1 from the following:

Theorem 4.2. *Let μ denote the μ -map associated to the Uhlenbeck compactification X of $\mathfrak{M}$. Then, using the complex orientation to identify $H^{2d}(X; \mathbb{Z}) \cong \mathbb{Z}$, we have, for all p with $-p \geq 2(4p_g + 2)$ and for all $\Sigma \in H_2(S)$,*

$$\mu(f)^m \cup \mu(\Sigma)^{d-m} = \begin{cases} 0, & \text{if } m > n \\ (d-n)!(2m_1 m_2)^{d-n} m_2(\Sigma \cdot \kappa_S)^{d-n}, & \text{if } m = n. \end{cases}$$

Here the factor $(2m_1 m_2)^{p_g(S)}$ appearing in Theorem 4.1 arises from the fact that $\mu(f)^n = (2m_1 m_2)^n \mu(\kappa)$ and that $d - 2n = p_g(S)$.

In order to prove Theorem 4.2, we shall introduce geometric divisors which will represent the cohomology class $\mu(f)$. Fix a smooth fiber f . Then there are exactly four line bundles θ of degree $m_1 m_2$ on f such that $\theta^{\otimes 2} = \mathcal{O}_f(\Delta)$. Each line bundle θ corresponds to a point of intersection of f with the branch divisor $B \subset \mathbb{F}$. If V is a rank two vector bundle over S with $c_1(V) = \Delta$, then $(V|f) \otimes (\theta)^{-1}$ is a vector bundle over f with trivial determinant. Thus by the Riemann-Roch theorem $\chi(f; (V|f) \otimes (\theta)^{-1}) = 0$.

For each integer c , fix a (Δ, c) -suitable line bundle L . Given an integer $b \leq c$, we define $\mathfrak{M}_b$ to be the moduli space of L -stable rank two bundles V with $c_1(V) = \Delta$ and $c_2(V) = b$. Given f and θ as above, define the divisor $\mathcal{Z}_b(f, \theta)$ as a set by:

$$\mathcal{Z}_b(f, \theta) = \{ V \in \mathfrak{M}_b : h^0(f; (V|f) \otimes (\theta)^{-1}) \neq 0 \}.$$

A calculation with the Grothendieck-Riemann-Roch theorem as in the proof of Proposition 1.1 in Chapter 5 of [4] show that we can use the divisors $\mathcal{Z}_b(f, \theta)$ to calculate the μ -map. More precisely, since the f_i are disjoint, suppose that, for all choices of $b > 0$ the intersection

$$\mathcal{Z}_{c-b}(f_{i_1}, \theta_{i_1}) \cap \cdots \cap \mathcal{Z}_{c-b}(f_{i_{n-b}}, \theta_{i_{n-b}}) = \emptyset,$$

that $\bigcap_{i=1}^n \mathcal{Z}_c(f_i, \theta_i) = J$ is compact and is contained in the Zariski open subset M of $\mathfrak{M}$, and that the $\mathcal{Z}_c(f_i, \theta_i)$ meet transversally along J . Here by (2.6), $\mathfrak{M}_{c-b} = \emptyset$ if $b > (-p-3)/4$ and so it is enough to check the above for $0 < b \leq (-p-3)/4$. In particular we must have $n > b$ for all $b \leq (-p-3)/4$. Then we can define $\mu([\Sigma])|J$ and arguments along the lines of the proof of Theorem 1.12 in Chapter 5 of [4] show that

$$\gamma_{w,p}(S)(f_1, \dots, f_n, [\Sigma], \dots, [\Sigma]) = (\mu([\Sigma])|J)^{d-n}.$$

Thus, Theorem 4.2 is a consequence of the following:

Theorem 4.3. *Suppose that $-p \geq 2(4p_g+2)$. Let $f_1, \dots, f_t$ denote distinct general fibers of π , and for each i choose θ_i a line bundle on f_i with $\theta_i^2 = \mathcal{O}_{f_i}(\Delta)$. Then:*

- (i) *For all $t \geq n$ and for all choices of $b > 0$, the intersection*

$$\mathcal{Z}_{c-b}(f_{i_1}, \theta_{i_1}) \cap \cdots \cap \mathcal{Z}_{c-b}(f_{i_{t-b}}, \theta_{i_{t-b}}) = \emptyset.$$

- (ii) *If $t > n$, then $\bigcap_{i=1}^t \mathcal{Z}_c(f_i, \theta_i) = \emptyset$, and moreover $\bigcap_{i=1}^n \mathcal{Z}_c(f_i, \theta_i)$ is compact and is contained in the Zariski open subset M of $\mathfrak{M}$. The intersection is transverse and is a fiber J of the map $M \rightarrow U \subseteq \mathbb{P}^n$.*
- (iii) *If Σ is a smooth curve in S , then the restriction of $\mu([\Sigma])$ to each of the m_2 connected components of J is equal to $(\Sigma \cdot f) \cdot [\Theta]$, where Θ is the theta divisor of the component.*

We begin by determining the set-theoretic intersection of the $\mathcal{Z}_c(f_i, \theta_i)$. Recall that we have associated to V the section A and the scheme Z on the double cover T . The following is straightforward:

Lemma 4.4. *Let $V \in \mathfrak{M}$. Then $V \in \mathcal{Z}_c(f, \theta)$ is and only if either the section A' of $\mathbb{F}$ corresponding to V meets B transversally at the point corresponding to θ or $\text{Supp } Z \cap \nu^{-1}(f) \neq \emptyset$. $\square$*

For the rest of the argument, we shall not worry about the case where the sections pass through the exceptional point, since this can be handled by symmetry. Arguing as in [4], Lemma 2.7 of Chapter 7, for a general choice of fibers $f_1, \dots, f_t$ and line bundles θ_i on f_i , and for all $s \leq r$, setting H_i to be the hyperplane of sections in $|\sigma + (N+s)f|$ passing through the point corresponding to θ_i , then $H_1 \cap \cdots \cap H_{N+2s+1} = \{A\}$ and the intersection of more than $N+2s+1$ of the H_i is empty. In particular, this means that the f_i are chosen so that the negative section σ does not meet the branch divisor at any of the f_i . Thus if V is a bundle whose associated section A is the negative section σ of $\mathbb{F}_N$ and V lies in the intersection $\bigcap_{i=1}^t \mathcal{Z}_c(f_i, \theta_i)$, then $\text{Supp } Z$ meets the preimage in T of f_i for all i .

A counting argument as in the proof of Proposition 2.9 of Chapter 7 in [4] then establishes (i) and (ii) of Theorem 4.3, at least set-theoretically, except possibly

for those V such that the corresponding section of $\mathbb{F}_N$ is the negative section. Likewise, arguments identical to the proof of (2.6) in Chapter 7 of [4] show that the intersection of the divisor $\mathcal{Z}_c(f_i, \theta_i)$ with the Zariski open subset M is reduced, so that the intersection in (ii) of Theorem 4.3 is transverse. Thus the only new case to consider is the possibility that the section associated to V is the negative section. We shall briefly outline the argument in this case; it is here that we must assume that $-p$ is sufficiently large. In particular, recalling that

$$n = \frac{d - p_g}{2} = \frac{-p - 4p_g - 3}{2},$$

and that we may assume that $b \leq (-p - 3)/4$, the condition $-p \geq 2(4p_g + 2)$ insures that

$$\begin{aligned} n - b &\geq n + \frac{p + 3}{4} = -\frac{p}{4} - \frac{8p_g + 3}{4} \\ &\geq \frac{8p_g + 4}{2} - \frac{8p_g + 3}{4} > 0. \end{aligned}$$

Thus the intersection in (i) is not over the empty collection of divisors $\mathcal{Z}_{c-b}(f_i, \theta_i)$, i.e. the intersection is always contained in a divisor $\mathcal{Z}_{c-b}(f_i, \theta_i)$.

Given t general fibers $f_1, \dots, f_t$, we can assume that they do not correspond to intersections of the negative section σ with $B + \mathfrak{d}_1 + \mathfrak{d}_2$. Now let V be a bundle whose associated section is σ . Suppose further that $c_2(V) = c - b$ and that V lies in the intersection $\mathcal{Z}_{c-b}(f_{i_1}, \theta_{i_1}) \cap \dots \cap \mathcal{Z}_{c-b}(f_{i_t-b}, \theta_{i_t-b})$ for $t \geq n$. We claim that, if $0 \leq b < (-p - 3)/4$, then $-p \leq 4p_g + 2N - 2$. It clearly suffices to assume that $t = n$. Let $\nu: T \rightarrow S$ be the double cover corresponding to σ . Then ν is not branched over the f_i . Writing ν^*V as an extension

$$0 \rightarrow \mathcal{O}_T(D) \rightarrow \nu^*V \rightarrow \mathcal{O}_T(\nu^*\Delta - D) \otimes I_Z \rightarrow 0,$$

we have by (2.5) that $-p = -p_1(\text{ad } V) + 4b = 4k - 6N + 1 + 2\ell(Z) + \delta + 4b$. Moreover $\text{Supp } Z$ must meet each of the two components of $\nu^{-1}(f_i)$ for $i = 1, \dots, t$. Thus $\ell(Z) \geq 2n - 2b$. But we also have

$$\begin{aligned} 2n &= d - p_g = -p - 4p_g - 3 \\ &= 4k - 6N + 1 + 2\ell(Z) + \delta - 4(k - N - 1) - 3 + 4b \\ &= -2N + 2 + 2\ell(Z) + \delta + 4b. \end{aligned}$$

Thus $2n \geq -2N + 2 + 4n - 4b + \delta + 4b$ and so $2n \leq 2N - 2$. This says that

$$-p \leq 4p_g + 2N - 2.$$

On the other hand, $2(4p_g + 2) = 4p_g + 4(k - N) \geq 4p_g + 4N > 4p_g + 2N - 2$. Thus with our assumptions on p there can be no bundle V in the intersection with an associated section equal to σ . Taking $b > 0$ establishes (i) in the case where the section associated to V is σ . Taking $b = 0$ shows that no bundle with associated section σ lies in the intersection $\bigcap_{i=1}^t \mathcal{Z}_c(f_i, \theta_i)$ if $t \geq n$. This establishes (i) and (ii).

Finally, to prove (iii) of Theorem 4.3, we must determine $\mu([\Sigma])|J$, where J is a fiber of the map $M \rightarrow U$. The argument again closely parallels that of [4] Chapter 7. A fiber of $M \rightarrow U$ determines and is determined by a generic double cover $T \rightarrow S$. There is a divisor D_0 on T such that every V in the fiber is of the form $\nu_* \mathcal{O}_T(D_0 + F) \otimes \rho^* \lambda$, for a line bundle $\lambda \in \text{Pic}^0 C$. Fix a smooth holomorphic multisection Σ of π , transverse to the double cover $T \rightarrow S$. Let Σ' be the inverse image of Σ under ν . There is a commutative diagram

$$\begin{array}{ccc} \Sigma' & \xrightarrow{\nu} & \Sigma \\ \rho \downarrow & & \downarrow \pi \\ C & \xrightarrow{g} & \mathbb{P}^1. \end{array}$$

Let $\mathcal{P}$ be the Poincaré bundle over $\text{Pic}^0 C \times C$. Let E be the divisor on Σ' induced by $D_0 + F$. Then there is a universal bundle over $\text{Pic}^0 C \times \Sigma$ of the form

$$(\text{Id} \times \nu)_* \left[\pi_2^* \mathcal{O}_{\Sigma'}(E) \otimes (\text{Id} \times \rho^*) \mathcal{P} \right].$$

Here $\pi_2: \text{Pic}^0 C \times \Sigma' \rightarrow \Sigma'$ is the second projection. The Chern classes of $\mathcal{V}_E$ only depend on the numerical equivalence class of E . Moreover, $p_1(\text{ad } \mathcal{V}_E) = p_1(\text{ad } \mathcal{V}_E \otimes q_2^* \lambda)$ for every line bundle λ on Σ , where $q_2: \text{Pic}^0 C \times \Sigma \rightarrow \Sigma$ is the projection. This has the effect of replacing $\mathcal{O}_{\Sigma'}(E)$ by $\mathcal{O}_{\Sigma'}(E) \otimes \nu^* \lambda$. Replacing Σ by 2Σ , replaces $\deg E$ by $2 \deg E$. Thus we can assume that $\deg E$ is even, and then after twisting by an appropriate λ we can assume that $\deg E = 0$. So as far as the Chern classes are concerned we may as well assume that $E = 0$, and we need to calculate

$$p_1(\text{ad}(\text{Id} \times \nu)_*(\text{Id} \times \rho)^* \mathcal{P}).$$

Now since cohomology commutes with flat base change, we have

$$(\text{Id} \times \nu)_*(\text{Id} \times \rho)^* \mathcal{P} = (\text{Id} \times \pi)^*(\text{Id} \times g)_* \mathcal{P}.$$

Thus we need to find

$$p_1(\text{ad}(\text{Id} \times \pi)^*(\text{Id} \times g)_* \mathcal{P}) = (\text{Id} \times \pi)^* p_1(\text{ad}(\text{Id} \times g)_* \mathcal{P}).$$

Let us first calculate $p_1(\text{ad}(\text{Id} \times g)_* \mathcal{P})$. A straightforward calculation using e.g. Lemma 2.14 of Chapter 7 of [4] shows the following

Lemma 4.5. *Let $f: X \rightarrow Y$ be a double cover with X and Y smooth, and let Υ be the line bundle on Y defining the double cover, so that $\Upsilon^{\otimes 2} = \mathcal{O}_Y(B)$, where B is the branch locus of f . If D is a divisor on X , then*

$$p_1(\text{ad } f_* \mathcal{O}_X(D)) = c_1(\Upsilon)^2 - (f_* D)^2 + 2f_*(D^2). \quad \square$$

Applying this in our situation, with $X = \text{Pic}^0 C \times C$ and $f = \text{Id} \times g$, we see that Υ is the pullback of a divisor on C and thus $c_1(\Upsilon)^2 = 0$. So we are left with calculating (in the sense of cycles) $(\text{Id} \times g)_* \mathcal{P}$ and $(\text{Id} \times g)_* [\mathcal{P}]^2$. Also, as in the proofs of (2.15) and (2.16) in Chapter 7 of [4], $(\text{Id} \times g)_* \mathcal{P} = 0$ and $[\mathcal{P}]^2 = -2r_1^*[\Theta] \cup r_2^*x$, where x

is the class of a point in C and Θ is the theta divisor on $\text{Pic}^0 C$, and r_1, r_2 are the first and second projections on $\text{Pic}^0 C \times C$. Thus

$$\begin{aligned} 2(\text{Id} \times g)_*[\mathcal{P}]^2 &= 2(\text{Id} \times g)_*(-2r_1^*[\Theta] \cup r_2^*x) \\ &= -4p_1^*[\Theta] \cup p_2^*y, \end{aligned}$$

where p_1, p_2 are the first and second projections on $\text{Pic}^0 C \times \mathbb{P}^1$ and y is the class of a point on $\mathbb{P}^1$. It follows that $-p_1(\text{ad } \mathcal{V}_E)/4 = q_1^*\Theta \cup q_2^*\pi^*y$.

Hence the slant product of $-p_1(\text{ad } \mathcal{V}_E)/4$ with $[\Sigma]$ is $(\deg \pi) \cdot [\Theta]$. Since $\deg \pi = (\Sigma \cdot f)$, we have now established (iii) of Theorem 4.3. This concludes the proof of Theorem 4.3 and thus of Theorem 4.1. $\square$

REFERENCES

1. S. K. Donaldson and P. B. Kronheimer, *The Geometry of Four-Manifolds*, Clarendon, Oxford, 1990.
2. R. Friedman, *Rank two vector bundles over regular elliptic surfaces*, Inventiones Math. **96** (1989), 283–332.
3. ———, *Vector bundles and $SO(3)$ -invariants for elliptic surfaces I* (to appear).
4. R. Friedman and J. W. Morgan, *Smooth 4-manifolds and complex surfaces* (to appear).

COLUMBIA UNIVERSITY, NEW YORK, NY 10027
E-mail address: `rf@math.columbia.edu`